



12th Oxford School on Neutron Scattering

September 2011

Introductory Theory

*Andrew Boothroyd
Oxford University*



Books on Neutron Scattering

- Theory

Lovesey, S.W. [formerly Marshall, W. and Lovesey, S.W.]

Theory of Neutron Scattering from Condensed Matter

O.U.P., 1984, 2 volumes, ~£25 each

Definitive formal treatment, but not for the faint-hearted!
(out of print)

Squires, G.L.

Intro. to the Theory of Thermal Neutron Scattering

C.U.P., 1978, Republished by Dover, 1996, ~£10.

More elementary than Lovesey, excellent for basic theory

Gunn, J.M.

Introductory Theory of Neutron Scattering

Lectures from the 1986 Oxford Summer School on Neutron Scattering
Adam Hilger, 1988.

Less formal approach, but not so comprehensive

- General (with introductory theory)

Willis, B.T.M. and Carlile, C.G.

Experimental Neutron Scattering,

O.U.P., 2009, £45

General introduction to all aspects of neutron scattering; v. good on methods

Furrer, A., Mesot, J., Strässle, T.

Neutron Scattering in Condensed Matter Physics

World Scientific, 2009, £24

Basic principles of neutron scattering and applications to a range of different materials and phenomena

Sivia, D.S.

Elementary Scattering Theory for X-ray and Neutron Users

O.U.P., 2011, £20

Basic principles of neutron scattering from a wave perspective

Methods of Experimental Physics, Vol. 23 A, B, C

Academic Press, 1987

25 separate review articles on theory, sources, instrumentation and science

Oxford Series on Neutron Scattering in Condensed Matter

O.U.P., 1988–2008

15 books on different aspects of neutron scattering

Scattering Experiments

- Beam of **radiation** (neutrons, photons, electrons, etc) incident on sample
- Measure **distribution** of radiation scattered from sample
- **Interaction potential** determines what is measured
- Radiation must be **coherent** (spatially or temporally or both)

Neutrons as Particles and Waves

Matter Wave:

- Oscillations → wave
Envelope → particle
- Particle separation $\gg \lambda$ - classical
“ “ $\ll \lambda$ - quantum (e.g. superfluid He)
- Increase Δ to define λ better:

Decrease Δ to define position better,
but lose information on λ .

Cannot define both Δ and λ to arbitrary precision
(Heisenberg Uncertainty Principle)

- Kinematics – Einstein, de Broglie

1) Energy: $E = hf$ $h = \text{Planck's constant}$
 $= \hbar\omega$ $f = \text{frequency}$
 $\hbar = h/2\pi, \omega = 2\pi f$

2) Momentum: $p = h/\lambda$ $\mathbf{k} = \text{wavevector}$
 $\mathbf{p} = \hbar\mathbf{k}$ $|\mathbf{k}| = 2\pi/\lambda$

Elastic Scattering from Bound Nuclei

Single nucleus

Weak disturbance of a plane wave

Result: plane wave + spherical wave

Model for neutrons interacting with a nucleus:

- Assumption - small fraction are scattered
- Justification - nuclear potential is short range
most neutrons 'miss' nucleus

Born
Approximation

[• Formal theory uses a *pseudopotential*: $V(\mathbf{r}) = (2\pi\hbar^2 b/m) \delta(\mathbf{r} - \mathbf{R})$]

Scattering from a line of nuclei

a) Normal incidence

What is the diffraction angle ?

For constructive interference
the path difference = λ

$$\sin 2\theta = \lambda/d$$

Suppose path difference = 2λ :

$$\sin 2\theta = 2\lambda/d$$

In general:

$$\sin 2\theta = n\lambda/d$$

b) Incident angle = diffracted angle

diffraction condition:

$$n\lambda = 2d \sin \theta$$

Notes

- At large distances, diffracted waves are plane waves
- N nuclei:
amplitude of diffracted wave $\sim N$
elsewhere, amplitude ~ 1

Elastic Scattering from a Crystal

a) Normal Incidence

In general, $AA' \neq AB$, so diffraction from 2nd column of atoms not usually in phase with diffraction from first.

Only achieve constructive interference when a and d are in special ratios.


b) Angle of incidence = angle of reflection

This time, $AA' = BB'$, so always achieve constructive interference from successive columns of atoms.


Hence, diffraction from a crystal occurs when

$$n\lambda = 2d \sin \theta$$

The Braggs — founders of crystallography



W.H. Bragg
(1862–1942)



W.L. Bragg
(1890–1971)

- Developed X-ray diffraction techniques for solving crystal structures (1913)
- Bragg's law:

$$n\lambda = 2d \sin \theta$$

Proceedings of the Cambridge Philosophical Society **17**, 43 (1914)

- Shared Nobel Prize (1915)

Debye-Waller factor

In reality, nuclei are not stationary :

Instantaneous positions

Time-average


- causes decrease in intensity of diffracted beam because waves are not so well in phase.
- Effect worsens as k ($=2\pi/\lambda$) and θ increase

- Bragg's Law the same, but $d \rightarrow \langle d \rangle$.
- smearing increases with temperature :

$$I = I_0 \exp\{-\langle(\mathbf{Q}\cdot\mathbf{u})^2\rangle\} = I_0 \exp(-2W)$$

Debye-Waller
Factor

Single crystal diffraction data from $\text{Nd}_{0.5}\text{Pb}_{0.5}\text{MnO}_3$ taken on the SXD diffractometer, courtesy of Dr Dave Keen (ISIS).



Particle Waves (again)

2 assumptions of quantum mechanics :

1. A particle is represented mathematically by a wavefunction, $\psi(\mathbf{r})$.
2. Probability of finding the particle in a (infinitesimal) volume dV is $|\psi(\mathbf{r})|^2 dV$.

Examples

(i) Infinite plane wave :

$$\begin{aligned}\psi &= \exp\{ikz\} \quad (= \cos kz + i \sin kz) \\ |\psi|^2 &= \psi \psi^* \\ &= \exp\{ikz\} \exp\{-ikz\} \\ &= 1\end{aligned}$$

→ 1 particle per unit volume everywhere

(ii) Spherical wave :

$$\begin{aligned}\psi &= -\frac{b \exp\{ikr\}}{r} \\ |\psi|^2 &= b^2/r^2\end{aligned}$$


→ density of particles falls off as $1/r^2$

Flux of particles


$$\begin{aligned}I &= \text{number incident normally on unit area per sec.} \\ &= \text{particle density} \times \text{velocity} \\ &= |\psi|^2 v \\ &= |\psi|^2 \hbar k / m\end{aligned}$$

Summary of Lecture 1

- Nucleus provides a weak perturbation to the incident neutrons, scattered neutrons are described by spherical waves:



- Diffraction from crystals:



d = spacing between planes
 θ = half the scattering angle

- Measure diffraction peaks \rightarrow d -spacings
 \rightarrow crystal structure
- Thermal motion of atoms does not affect use of Bragg's Law, but does reduce peak intensities from their values for a perfectly rigid structure.

Cross-Sections

Total cross-section

Total cross-section σ is defined by,

$$\sigma = \frac{\text{total no. particles scattered in all directions per sec.}}{\text{incident flux } (I_0)}$$

(i) Classical case – scattering from a solid sphere, radius a

$$\text{No. particles scattered per sec.} = I_0 \times \pi a^2$$

$$\rightarrow \sigma = \pi a^2$$

(ii) Quantum case – scattering from an isolated nucleus

$$\begin{aligned} \text{Incident wave,} & \quad \psi_0 = \exp \{ ikz \} \\ \text{Incident flux,} & \quad I_0 = |\psi_0|^2 v = v \end{aligned}$$

$$\text{Scattered wave,} \quad \psi^{\text{sc}} = -\frac{b \exp \{ ikr \}}{r}$$

$$\begin{aligned} \text{Scattered flux,} & \quad I^{\text{sc}} = |\psi^{\text{sc}}|^2 v = b^2 v / r^2 \\ & \text{at distance } r \end{aligned}$$

$$\begin{aligned} \text{Total no. particles scattered per sec.} & = I^{\text{sc}} \times \text{total area} \\ & = b^2 v / r^2 \times 4\pi r^2 \\ & = 4\pi b^2 v \end{aligned}$$

$$\rightarrow \sigma = 4\pi b^2$$

Notes:

- σ is the *effective area* of the target as viewed by the incident neutrons
- if the target is a nucleus, then b is the **nuclear scattering length**; b is the effective range of the nuclear potential
- units of b : Fermi (f) 1 Fermi = 10^{-15} m
 " " σ : barn (b) 1 barn = 10^{-28} m²

Differential cross-section

Differential cross-section, $\frac{d\sigma}{d\Omega}$ is defined by,

$$\frac{d\sigma}{d\Omega} = \frac{\text{No. particles scattered into solid angle } d\Omega \text{ per sec.}}{I_0 \times d\Omega}$$

Solid angle subtended by detector at sample is $\Delta\Omega = A/L^2$

From definition of $\frac{d\sigma}{d\Omega}$, no. particles detected per sec. = $I_0 \Delta\Omega \frac{d\sigma}{d\Omega}$

but also, " " " " " = $|\psi^{sc}|^2 v \times A$

$$\rightarrow \frac{d\sigma}{d\Omega} = \frac{|\psi^{sc}|^2}{|\psi_0|^2} L^2$$

Example: isolated nucleus

At detector scattered wave is $\psi^{sc} = -\frac{b \exp\{ikL\}}{L}$

$$\rightarrow \frac{d\sigma}{d\Omega} = b^2 = \frac{\sigma}{4\pi}$$

Note:

- units of $\frac{d\sigma}{d\Omega}$: barns (steradian)⁻¹ (b sr⁻¹)

Scattering cross-section for an assembly of nuclei

Recall :
$$\frac{d\sigma}{d\Omega} = \frac{|\psi^{sc}|^2}{|\psi_0|^2} L^2$$

At detector,

$$\begin{aligned}\psi_0^{sc} &= -\frac{b_0 \exp\{ikL\}}{L} \\ \psi_n^{sc} &= -\frac{b_n \exp\{ik(L+\Delta L_n)\}}{(L+\Delta L_n)}\end{aligned}$$

What is ΔL_n ?

$$\begin{aligned}\Delta L_n &= An + nB \\ &= \frac{\mathbf{k}_i \cdot \mathbf{r}_n}{k} - \frac{\mathbf{k}_f \cdot \mathbf{r}_n}{k}\end{aligned}$$

$$\begin{aligned}\rightarrow k\Delta L_n &= (\mathbf{k}_i - \mathbf{k}_f) \cdot \mathbf{r}_n \\ &= \mathbf{Q} \cdot \mathbf{r}_n\end{aligned}$$

Total scattered wave,

$$\psi^{sc} = \sum_n \psi_n^{sc} = -\frac{\exp\{ikL\}}{L} \sum_n b_n \exp\{i\mathbf{Q} \cdot \mathbf{r}_n\}$$

Cross-section :

$$\frac{d\sigma}{d\Omega} = \left| \sum_n b_n \exp\{i\mathbf{Q} \cdot \mathbf{r}_n\} \right|^2$$

Bragg diffraction from a rigid crystal

Crystal is a periodic array of atoms.

Lattice is a periodic array of points representing the periodicity of the crystal. The lattice points are displaced from the origin by lattice vectors

$$\mathbf{l} = n_1\mathbf{a} + n_2\mathbf{b} + n_3\mathbf{c}, \quad (n_1, n_2, n_3 \text{ integers})$$

Unit cell is a building block from which the crystal is constructed.

Usually it is a parallelepiped with edges $\mathbf{a}, \mathbf{b}, \mathbf{c}$:

Cross-section :

$$\frac{d\sigma}{d\Omega} = \left| \sum_n b_n \exp \{ i\mathbf{Q} \cdot \mathbf{r}_n \} \right|^2$$

Position of nucleus \mathbf{r}_n :

$$\mathbf{r}_n = \mathbf{l} + \mathbf{d}$$

$$\rightarrow \frac{d\sigma}{d\Omega} = \left| \sum_{\mathbf{l}} \exp \{ i\mathbf{Q} \cdot \mathbf{l} \} \sum_{\mathbf{d}} b_{\mathbf{d}} \exp \{ i\mathbf{Q} \cdot \mathbf{d} \} \right|^2$$

Coherent (Bragg) scattering occurs when all terms in \mathbf{l} sum are equal, i.e.

$$\exp \{ i\mathbf{Q} \cdot \mathbf{l} \} = 1 \quad \text{for all } \mathbf{l}$$

Which values of \mathbf{Q} satisfy this equation? Answer:

$$\mathbf{Q} = h\mathbf{a}^* + k\mathbf{b}^* + l\mathbf{c}^* \quad (h, k, l \text{ integers})$$

where,

$$\mathbf{a}^* = (2\pi/v_0) \mathbf{b} \times \mathbf{c}, \quad \mathbf{b}^* = (2\pi/v_0) \mathbf{c} \times \mathbf{a}, \quad \mathbf{c}^* = (2\pi/v_0) \mathbf{a} \times \mathbf{b}$$

and $v_0 = \mathbf{a} \cdot \mathbf{b} \times \mathbf{c} = \mathbf{b} \cdot \mathbf{c} \times \mathbf{a} = \mathbf{c} \cdot \mathbf{a} \times \mathbf{b}$.

Note also that $\mathbf{a} \cdot \mathbf{a}^*$ etc = 2π , and $\mathbf{a} \cdot \mathbf{b}^*$ etc = 0

Now consider summation over position vector \mathbf{d} .

Write \mathbf{d} in terms of fractional coordinates $(x_{\mathbf{d}}, y_{\mathbf{d}}, z_{\mathbf{d}})$ of nucleus

$$\mathbf{d} = x_{\mathbf{d}}\mathbf{a} + y_{\mathbf{d}}\mathbf{b} + z_{\mathbf{d}}\mathbf{c}$$

When \mathbf{Q} satisfies the condition $\mathbf{Q} = h\mathbf{a}^* + k\mathbf{b}^* + l\mathbf{c}^*$, then

$$\begin{aligned}\frac{d\sigma}{d\Omega} &= N^2 \left| \sum_{\mathbf{d}} b_{\mathbf{d}} \exp \{ i (h\mathbf{a}^* + k\mathbf{b}^* + l\mathbf{c}^*) \cdot (x_{\mathbf{d}}\mathbf{a} + y_{\mathbf{d}}\mathbf{b} + z_{\mathbf{d}}\mathbf{c}) \} \right|^2 \\ &= N^2 |F_{hkl}|^2 \quad (N \text{ is the no. unit cells in the crystal})\end{aligned}$$

where,

$$F_{hkl} = \sum_{\mathbf{d}} b_{\mathbf{d}} \exp \{ 2\pi i (hx_{\mathbf{d}} + ky_{\mathbf{d}} + lz_{\mathbf{d}}) \}$$

F_{hkl} is known as the **structure factor** for the reflection (hkl) .

Reciprocal Lattice

Peter Paul Ewald
(1888–1985)
The inventor of the
reciprocal lattice



Strong elastic scattering occurs when

$$\mathbf{Q} = \boldsymbol{\tau}_{hkl} \quad (\text{Laue condition})$$

where, $\boldsymbol{\tau}_{hkl} = h\mathbf{a}^* + k\mathbf{b}^* + l\mathbf{c}^*$

The set of all vectors $\{\boldsymbol{\tau}_{hkl}\}$ is called the **Reciprocal Lattice**.



Max von Laue
(1879–1960)
Nobel Prize (1914)

2 properties:

- (i) $\boldsymbol{\tau}_{hkl}$ is normal to the plane (hkl) .
- (ii) $|\boldsymbol{\tau}_{hkl}| = 2\pi/d_{hkl}$

Bragg \equiv Laue:

$$\begin{aligned} |\mathbf{Q}| &= |\boldsymbol{\tau}_{hkl}| && \text{Laue Condition} \\ \rightarrow \frac{4\pi}{\lambda} \sin \theta &= 2\pi/d_{hkl} \\ \rightarrow \lambda &= 2d \sin \theta && \text{Bragg's Law} \end{aligned}$$

Summary of Lecture 2

- σ = total scattering cross-section
– probability that the neutron is scattered
- $\frac{d\sigma}{d\Omega}$ = differential scattering cross-section
– probability that the neutron is scattered into a specified direction

- For elastic scattering from a rigid structure

$$\frac{d\sigma}{d\Omega} = \left| \sum_n b_n \exp \{ i\mathbf{Q} \cdot \mathbf{r}_n \} \right|^2$$

- For a rigid crystal, Bragg scattering occurs when

$$\mathbf{Q} = \boldsymbol{\tau}_{hkl} \quad (\text{Laue condition})$$

where,

$$\boldsymbol{\tau}_{hkl} = h\mathbf{a}^* + k\mathbf{b}^* + l\mathbf{c}^* \quad (\text{reciprocal lattice vectors})$$

The cross-section for Bragg scattering is given by

$$\frac{d\sigma}{d\Omega} = N^2 |F_{hkl}|^2$$

where,

$$F_{hkl} = \sum_{\mathbf{d}} b_{\mathbf{d}} \exp \{ 2\pi i (hx_{\mathbf{d}} + ky_{\mathbf{d}} + lz_{\mathbf{d}}) \} \quad (\text{structure factor})$$

- Corollary: for a non-rigid crystal:

$$F_{hkl} = \sum_{\mathbf{d}} \exp(-W_{\mathbf{d}}) b_{\mathbf{d}} \exp \{ 2\pi i (hx_{\mathbf{d}} + ky_{\mathbf{d}} + lz_{\mathbf{d}}) \}$$

Coherent and Incoherent Scattering

$$\frac{d\sigma}{d\Omega} = \left| \sum_n b_n \exp \{ i\mathbf{Q} \cdot \mathbf{r}_n \} \right|^2$$

Recall: b_n characterizes the range of the neutron-nucleus interaction.

b_n depends upon :

- (i) which element;
- (ii) which isotope;
- (iii) relative spins of neutron and nucleus.

In principle, we can calculate $\frac{d\sigma}{d\Omega}$ exactly if we know the isotope and spin state of every nucleus. Not feasible in practice.

Simplifying assumption

Assume that distribution of isotopes and spin states is **random** and **uncorrelated** between the sites.

→ $\frac{d\sigma}{d\Omega}$ for one particular sample is the same as the **average** over many samples with same nuclear positions

$$\rightarrow \frac{d\sigma}{d\Omega} \approx \overline{\frac{d\sigma}{d\Omega}} \quad \text{ensemble average}$$

In order to proceed we need \bar{b} and $\overline{b^2}$
(for formulae see notes in section B of tutorial problems).

Ensemble averaging

Suppose sample contains only 1 type of atom, which has 3 different isotopes:

isotope	natural abundance	scattering length
●	50 %	b_B
●	25 %	b_R
●	25 %	b_G

$$\bar{b} = 0.5 b_B + 0.25 b_R + 0.25 b_G$$

$$\overline{b^2} = 0.5 b_B^2 + 0.25 b_R^2 + 0.25 b_G^2$$

Note that,

$$\frac{d\sigma}{d\Omega} = \left| \sum_n b_n \exp \{ i\mathbf{Q} \cdot \mathbf{r}_n \} \right|^2$$

$$= \sum_n \sum_m b_n b_m \exp \{ i\mathbf{Q} \cdot (\mathbf{r}_n - \mathbf{r}_m) \}$$

Ensemble averaging \rightarrow replace $b_n b_m$ by $\overline{b_n b_m}$

Sites uncorrelated \rightarrow

$$\overline{b_n b_m} = \overline{b}_n \overline{b}_m \quad \text{if } n \neq m$$

$$= \overline{b}_n^2 \quad \text{if } n = m$$

Therefore,

$$\frac{d\sigma}{d\Omega} = \sum_{n \neq m} \sum_m \overline{b}_n \overline{b}_m \exp \{ i\mathbf{Q} \cdot (\mathbf{r}_n - \mathbf{r}_m) \} + \sum_{n=m} \overline{b}_n^2$$

$$= \sum_n \sum_m \overline{b}_n \overline{b}_m \exp \{ i\mathbf{Q} \cdot (\mathbf{r}_n - \mathbf{r}_m) \} + \sum_{n=m} (\overline{b}_n^2 - \overline{b}_n^2)$$

coherent scattering
incoherent scattering

coherent scattering — correlations between the same, and different nuclei — **interference, structure** (also collective dynamics)

incoherent scattering — no information on structure — **'flat background'** (also dynamics of single particles)

Values of \overline{b} and \overline{b}^2 are tabulated (e.g. *Neutron News* vol. 3 No. 3 (1992) pp29-37 and <http://www.ncnr.nist.gov/resources/n-lengths/>)

Often written as

$$\sigma_{\text{coh}} = 4\pi \overline{b}^2$$

and

$$\sigma_{\text{inc}} = 4\pi (\overline{b}^2 - \overline{b}^2)$$

Examples

	σ_{coh} (barns)	σ_{inc} (barns)
hydrogen	1.8	80.2
carbon	5.6	0
vanadium	0	5

Examples of coherent and incoherent scattering

(i) Bragg diffraction from a powdered crystal

(i) Elastic scattering from a liquid or glass

Magnetic Scattering

- Neutron is uncharged, but possesses a magnetic dipole moment μ_n ($\sim 0.001\mu_B$) which can interact with magnetic fields from unpaired electrons via :

- (i) the intrinsic spin dipole moment of the electron,
- (ii) magnetic fields produced by orbital motion of electrons.

- Strength of magnetic interaction: $\sigma_{\text{mag}} \sim r_0^2 \sim 0.1$ barn
" " nuclear " $\sigma_{\text{coh}} \sim b^2 \sim 1$ barn

so similar magnitude.

- Theory similar to nuclear scattering except scatter from magnetic moments in sample, and this occurs via a **vector interaction**

$$V_M(\mathbf{r}) = -\mu_n \cdot \mathbf{B}(\mathbf{r})$$

- Neutron probes component of the atomic moment perpendicular to \mathbf{Q} .


- Neutrons scatter from electrons in atomic orbitals :
Smearred out in space

→ weaker scattering at higher angles
(like Debye-Waller factor)

Intensity fall-off described by a **magnetic form factor**
(similar to atomic form factor used in x-ray diffraction)

Diffraction from a Magnetic Structure

1. Ferromagnet




$$I_M \propto \sin^2\theta |F_M|^2 \quad (\theta \text{ is angle between } \mu \text{ and } Q)$$

where,

$$F_M = f(Q) \sum_j \mu_j \exp(i\mathbf{Q} \cdot \mathbf{r}_j) \quad \text{Magnetic structure factor}$$

2. Antiferromagnet



Neutron Polarization

- Neutron has spin 1/2, so moment is \uparrow or \downarrow relative to a magnetic field. Can have different scattering cross-sections according to the neutron spin state before and after scattering:

$$\begin{array}{ccc}
 \uparrow & \rightarrow & \uparrow \\
 \uparrow & \rightarrow & \downarrow \\
 \downarrow & \rightarrow & \uparrow \\
 \downarrow & \rightarrow & \downarrow \\
 \text{[or } & \mathbf{P}_i & \rightarrow & \mathbf{P}_f \text{ if initial and final field directions different]}
 \end{array}$$

→ polarization analysis

- Torque on magnetic dipole moment in magnetic field \mathbf{B} is

$$\mathbf{T} = \boldsymbol{\mu} \times \mathbf{B}$$

Eq. of motion:

Torque = rate of change of angular momentum

and angular momentum $\propto \boldsymbol{\mu}$

$$\rightarrow \frac{d\boldsymbol{\mu}}{dt} \propto \boldsymbol{\mu} \times \mathbf{B}$$

Consider 2 cases :

(i) $\boldsymbol{\mu}$ parallel to \mathbf{B}

no change in neutron spin state ('non- spin-flip scattering')

(ii) $\boldsymbol{\mu}$ perpendicular to \mathbf{B}

neutron spin precesses in field ('spin-flip scattering')

Neutron Inelastic Scattering

Kinematics (again)

Scattering triangle ($\mathbf{k}_i \neq \mathbf{k}_f$):

\mathbf{k}_i = incident wavevector

\mathbf{k}_f = final scattered wavevector

\mathbf{Q} = scattering vector

- Momentum transfer $\hbar\mathbf{Q} = \hbar(\mathbf{k}_i - \mathbf{k}_f)$
- Energy transfer $\hbar\omega = E_i - E_f = \frac{\hbar^2}{2m} (k_i^2 - k_f^2)$

A scattering event is characterised by (\mathbf{Q}, ω)

Accessible region of (\mathbf{Q}, ω) space :

Neutron Cross-Section

Suppose detector can analyse energy of neutrons.

Define the *double differential scattering cross-section* :

$$\frac{d^2\sigma}{d\Omega dE_f} = \frac{\text{No. particles scattered per sec. into solid angle } d\Omega \text{ with final energies between } E_f \text{ and } E_f + dE_f}{I_0 \times d\Omega \times dE_f}$$

Numerator depends implicitly on 5 factors :

- (i) $d\Omega$
- (ii) dE_f
- (iii) speed of scattered neutrons, $v_f = \hbar k_f / m$
- (iv) density of incident neutrons $|\psi_0|^2$
- (v) $S(\mathbf{Q}, \omega)$, the probability that system can change its energy by an amount $\hbar\omega$, accompanied by a momentum change $\hbar\mathbf{Q}$

In denominator, remember $I_0 = |\psi_0|^2 v_i = |\psi_0|^2 \hbar k_i / m$

Hence, these factors together give

$$\frac{d^2\sigma}{d\Omega dE_f} = \frac{k_f}{k_i} S(\mathbf{Q}, \omega)$$

Notes :

- $S(\mathbf{Q}, \omega)$ contains all the physics of the system
 - scattering function / response function
- the k_f/k_i factor is sometimes important, for example if the neutron loses a lot of energy ($k_f \ll k_i$) then the intensity is much reduced.

Scattering from lattice vibrations in a crystal

(Example of coherent inelastic scattering)

Phonon - quantum of lattice vibrational energy

Consider 1-*d* chain of identical atoms:

(1) Transverse vibrational mode

(2) Longitudinal vibrational mode (sound wave)

- Equivalent wavevectors

In general,

$$\mathbf{k}_{\text{ph}} \equiv \mathbf{k}_{\text{ph}} + \boldsymbol{\tau}$$

- Phonon dispersion curve

Energy $\hbar\omega_{\text{ph}}$ of a phonon depends on \mathbf{k}_{ph}

- Scattering from phonons

Peaks occur when

$$\begin{cases} \hbar\omega = \hbar\omega_{\text{ph}} \\ \hbar\mathbf{Q} = \hbar(\mathbf{k}_{\text{ph}} + \boldsymbol{\tau}) \end{cases}$$

(1) Longitudinal :

(2) Transverse :

- Inelastic scattering cross-section for phonons

Consider a *static* sinusoidal distortion of the lattice:

Position of n^{th} atom $x_n = nd + \alpha \sin(k_{\text{ph}}nd)$

Elastic scattering cross-section :

$$\begin{aligned} \frac{d\sigma}{d\Omega} &= \left| \sum_n b_n \exp \{ i\mathbf{Q} \cdot \mathbf{r}_n \} \right|^2 \\ &= \left| \sum_n b \exp \{ iQ(nd + \alpha \sin(k_{\text{ph}}nd)) \} \right|^2 \end{aligned}$$

Can make Taylor expansion in $Q\alpha$ when $Q\alpha \ll 1$:

$$\rightarrow \frac{d\sigma}{d\Omega} = \left| \sum_n b \left[\underbrace{1}_{(1)} + \underbrace{iQ\alpha \sin(k_{\text{ph}}nd)}_{(2)} + \dots \right] \exp \{ iQnd \} \right|^2$$

1st term (1)

\rightarrow Bragg peak at $Q = m (2\pi/d)$ ($m = \text{integer}$)
Intensity $\propto b^2$

2nd term (2): write $\sin x = (e^{ix} - e^{-ix})/2i$

$$\rightarrow \left| \sum_n bQ\alpha \left[\exp \{ i(Q + k_{\text{ph}})nd \} - \exp \{ i(Q - k_{\text{ph}})nd \} \right] \right|^2$$

\rightarrow peaks at $Q = m (2\pi/d) \pm k_{\text{ph}}$
Intensity $\propto b^2Q^2\alpha^2$

Lattice vibration – dynamic, sinusoidal distortion of the lattice

Inelastic scattering cross-section as for static case but conserve energy as well

$$\rightarrow \text{Peaks in } \frac{d^2\sigma}{d\Omega dE_f} \text{ when } \begin{cases} \hbar\omega = \pm\hbar\omega_{\text{ph}} \\ \hbar\mathbf{Q} = \hbar(\boldsymbol{\tau} \pm \mathbf{k}_{\text{ph}}) \end{cases}$$

$$\text{Intensity} \propto b^2 Q^2 \alpha^2$$

$$\propto \frac{b^2 Q^2}{\omega_{\text{ph}}} \quad (\alpha^2 \propto 1/\omega_{\text{ph}})$$

Spin Waves

Ground state of ferromagnet:

Displace one spin:

Displacement propagates through lattice as wave with wavevector k_{mag}

Magnon dispersion curve :

Notes

- Angular momentum (spin) of the crystal is reduced by 1 unit (of \hbar)
 - spin of neutron changes by 1 unit to conserve angular momentum
 - spin flip scattering
- Intensity varies with magnetic form factor - decreases with $|\mathbf{Q}|$.

Summary of Coherent Inelastic Scattering

- Double differential scattering cross-section :

$$\frac{d^2\sigma}{d\Omega dE_f} = \frac{k_f}{k_i} S(\mathbf{Q}, \omega)$$

- Propagating excitations (e.g. lattice vibs., spin waves) $S(\mathbf{Q}, \omega)$ has peaks

$$\text{when } \begin{cases} \hbar\omega = \pm\hbar\omega_{\text{ph}} \\ \hbar\mathbf{Q} = \hbar(\boldsymbol{\tau} \pm \mathbf{k}_{\text{ph}}) \end{cases}$$

- The size of the peaks in $S(\mathbf{Q}, \omega)$ varies according to

(i) Phonons

$$S(\mathbf{Q}, \omega) \propto \exp\{-2W(\mathbf{Q}, T)\} \times |\mathbf{G}(\mathbf{Q})|^2 \times [n(\omega_{\text{ph}}) + 1] \times \frac{1}{\omega_{\text{ph}}} \times Q^2$$

(ii) Spin waves

$$S(\mathbf{Q}, \omega) \propto \exp\{-2W(\mathbf{Q}, T)\} \times [n(\omega_{\text{mag}}) + 1] \times \frac{1}{\omega_{\text{mag}}} \times f^2(Q)$$

- Excitations can be measured in neutron energy loss or neutron energy gain, but remember that $S(\mathbf{Q}, \omega)$ has the property,

$$S(\mathbf{Q}, -\omega) = \exp(-\hbar\omega/k_B T) \times S(\mathbf{Q}, \omega)$$