

# Neutron Sources

Oxford School on Neutron Scattering  
8<sup>th</sup> September 2015


Ken Andersen

# Summary


- Neutron facilities
  - history, overview & trends
- Reactor-based sources
  - Institut Laue-Langevin
- Short-pulse spallation sources
  - ISIS
- Components of a pulsed spallation neutron source
  - accelerator
  - target
  - moderators
- Neutron source time structure
  - the time of flight method
- Long-pulse neutron sources




# The first neutron source



James Chadwick:  
used Polonium as alpha emitter on Beryllium




# Evolution of neutron sources




(Updated from *Neutron Scattering*, K. Sköld and D. L. Price, eds., Academic Press, 1986)

# Nuclear Fission




# Evolution of neutron sources




(Updated from *Neutron Scattering*, K. Sköld and D. L. Price, eds., Academic Press, 1986)

# Evolution of neutron sources




(Updated from *Neutron Scattering*, K. Sköld and D. L. Price, eds., Academic Press, 1986)

# Nuclear Spallation




# Evolution of neutron sources



(Updated from *Neutron Scattering*, K. Sköld and D. L. Price, eds., Academic Press, 1986)

# Evolution of neutron sources



(Updated from *Neutron Scattering*, K. Sköld and D. L. Price, eds., Academic Press, 1986)




# Slow Neutrons vs Light

	light	neutrons
$\lambda$	$< \mu\text{m}$	$< \text{nm}$
E	$> \text{eV}$	$> \text{meV}$
penetration	$\sim \mu\text{m}$	$\sim \text{cm}$
$\theta_c$	$90^\circ$	$1^\circ$
B	$10^{18}$ p/cm <sup>2</sup> /ster/s (60W lightbulb)	$10^{14}$ n/cm <sup>2</sup> /ster/s (60MW reactor)
spin	1	$\frac{1}{2}$
interaction	electromagnetic	strong force, magnetic
charge	0	0

# Why neutrons?

- Thermal neutrons have wavelengths similar to inter-atomic distances
- Thermal neutrons have energies comparable to lattice vibrations
- Neutrons are non-destructive
- Neutrons interact weakly
  - they penetrate into the bulk
- Neutrons interact via a simple point-like potential
  - amplitudes are straightforward to interpret
- Neutrons have a magnetic moment
  - great for magnetism
- Neutrons see a completely different contrast to x-rays
  - e.g. hydrogen is very visible

# Why neutrons?




X-rays  
Neutrons

Element (Z)

# Main European neutron sources 2017




# Major neutron sources in the world





# ILL Reactor Neutron Source











# ILL Reactor Neutron Source






# ILL Reactor Neutron Source



# ILL Reactor Neutron Source


- Highly-enriched uranium
- Compact design for high brightness
- Heavy-water cooling
- Single control rod
- 57MW thermal power
- Cold, thermal, hot sources




# ILL Reactor Neutron Source

- Highly-enriched uranium
- Compact design for high brightness
- Heavy-water cooling
- Single control rod
- 57MW thermal power
- Cold, thermal, hot sources

	cold	thermal	hot
moderator	liquid D <sub>2</sub>	Liquid D <sub>2</sub> O	graphite
moderator temperature	20K	300K	2000K
neutron wavelength	3→20Å	1→3Å	0.3→1Å




# ILL Reactor Neutron Source







# ILL Reactor Neutron Source



# ILL Reactor Neutron Source



# ILL Moderator Brightnesses




# Spallation vs Fission

## Fission

200 MeV/fission

$2.35 - 1 = 1.35$  neutrons freed

=> 150 MeV/neutron







# Spallation vs Fission

## Fission

200 MeV/fission  
 $2.35 - 1 = 1.35$  neutrons freed  
 $\Rightarrow 150$  MeV/neutron



## Spallation




1 GeV proton in:  
250 MeV becomes mass (endothermic reaction)  
30 neutrons freed  
 $\Rightarrow 25$  MeV/neutron


# Spallation vs Fission

## Fission

200 MeV/fission  
 $2.35 - 1 = 1.35$  neutrons freed  
 $\Rightarrow 150$  MeV/neutron



## Spallation



1 GeV proton in:  
250 MeV becomes mass (endothermic reaction)  
30 neutrons freed  
 $\Rightarrow 25$  MeV/neutron

6x more neutrons per unit heat



# Spallation Sources

- Spallation: 10x higher neutron brightness per unit heat
  - about 6x more neutrons per unit heat
  - about ½ the production volume
- 1 MW spallation source = 10 MW reactor
  - e.g. 800 MeV at 1.25 mA (PSI)
  - e.g. 3 GeV at 0.4 mA (J-PARC)
- Peak brightness >> time-average brightness

# Spallation Sources

- Spallation: 10x higher neutron brightness per unit heat
  - about 6x more neutrons per unit heat
  - about ½ the production volume
- 1 MW spallation source = 10 MW reactor
  - e.g. 800 MeV at 1.25 mA (PSI)
  - e.g. 3 GeV at 0.4 mA (J-PARC)

- Peak brightness  $\gg$  time-average brightness



# De Broglie Relations

Particle	Wave
$p = mv$	$p = \hbar k = h/\lambda$
$E = \frac{1}{2}mv^2$	$E = \hbar\omega = hf$

$$\hbar = h/2\pi$$

$$h = 6.6 \times 10^{-34} \text{ J} \cdot \text{s}$$

$$m_n = 1.67 \times 10^{-27} \text{ kg}$$

$$\lambda = h / mv$$


$$\lambda[\text{\AA}] = 3.956 / v[\text{m/ms}]$$

$$t[\text{ms}] = L[\text{m}] \times \lambda[\text{\AA}] / 3.956$$

# The Time-of-Flight (TOF) Method


$$t[\text{ms}] = L[\text{m}] \times \lambda[\text{\AA}] / 3.956$$

distance





# ISIS, UK (160kW)



Science & Technology Facilities Council

# ISIS



# SNS, Oak Ridge, USA (1MW)






# J-PARC, Tokai, Japan (500kW)



# J-PARC, Tokai, Japan (500kW)





# ESS, Lund, Sweden (5MW in 2025)



# Short-Pulse Spallation Sources


- Accelerator
  - $H^-$  ion source
  - Linear accelerator (“linac”)
  - Stripper to convert  $H^-$  to  $H^+$
  - Synchrotron
- Target
- Reflector
- Moderators




# Linear accelerator: LINAC



# Linear accelerator: LINAC




# SNS ion source: H<sup>-</sup>






# Different types of Linac

Drift-Tube Linac (DTL)




Radio-Frequency Quadrupole (RFQ)

Elliptical cavities




“Low  $\beta$ ” / “High  $\beta$ ”

$$\beta = v/c$$




# Synchrotron

- Synchronise:
  - B-field: bend
  - E-field: accelerate
  - E & B field: focus
  - magnets to each other
- Injection
  - stripper foil
- Extraction
  - kicker magnet



# Synchrotron


- Synchronise:
  - B-field: bend
  - E-field: accelerate
  - E & B field: focus
  - magnets to each other
- Injection
  - stripper foil
- Extraction
  - kicker magnet








# ISIS target 1: solid tungsten







# SNS Target Configuration




# SNS target: liquid mercury



# ESS target



# ISIS TS2 Target




# Target-Reflector-Moderator Neutronics


- Target produces neutrons in  $> \text{MeV}$  range
- Moderators contain H to thermalise neutrons
  - largest scattering cross-section (80b)
  - lower mass: same as neutron
  - on average,  $\frac{1}{2}$  energy lost per collision
  - 100 MeV  $\rightarrow$  10 meV requires about 25 collisions
- Moderators embedded in reflector, usually  $\text{D}_2\text{O}$ -cooled Be
  - minimal absorption
  - large scattering cross-section (8b)
  - little thermalisation




# Target-Reflector-Moderator Neutronics




# Target-Reflector-Moderator Neutronics




# Target-Reflector-Moderator Neutronics



# Target-Reflector-Moderator Neutronics




# Target-Reflector-Moderator Neutronics






# Target-Reflector-Moderator Neutronics







# Target-Reflector-Moderator Neutronics




# Target-Reflector-Moderator Neutronics

Decoupling  
Poisoning



# Time-of-flight (TOF) resolution

$$t[\text{ms}] = L[\text{m}] \times \lambda[\text{\AA}] / 3.956$$






# Time-of-flight (TOF) resolution



$$t[\text{ms}] = L[\text{m}] \times \lambda[\text{\AA}] / 3.956$$

$$\Rightarrow \Delta\lambda[\text{\AA}] = \Delta t[\text{ms}] \times 3.956 / L[\text{m}]$$





# Moderator Decoupling and Poisoning


## J-PARC H<sub>2</sub> moderators at 1MW




# Moderator Decoupling and Poisoning



# SNS moderators




# ISIS TS2 Target





# Moderator Temperature

ISIS-TS1 moderators at 160kW



# Beyond Short-Pulse Limits



SNS instantaneous power on target:

17kJ in 1 $\mu$ s: 17 x

Reaches limits of spallation source technology:  
 shock waves in target, space charge density in  
 accelerator ring, ...





# Beyond Short-Pulse Limits



SNS instantaneous power on target:


17kJ in 1 $\mu$ s: 17 x

ESS instantaneous power on target: 125MW

360kJ in 2.86ms




# Long-pulse performance



# Adapting the pulse width


## Short-Pulse Source

- set pulse width by choosing moderator



## Long-Pulse Source (ESS)

- set pulse width using pulse-shaping chopper





# Summary

- Neutron facilities
  - overview & trends
- Reactor-based sources
  - Institut Laue-Langevin
- Fission vs Spallation
  - ISIS
- Components of a pulsed spallation neutron source
  - accelerator
  - target
  - moderators
- Neutron source time structure
  - the time of flight method
- Long-pulse neutron sources

# Thank You!

